

Endoscopy Department

ERCP (Endoscopic Retrograde Cholangio Pancreatography)

Important Information

Before your appointment

- If you are taking **Warfarin** please read the '[Alert for patients on Warfarin](#)' carefully as you may need to have an INR test 7 days before.
- All other medication should be taken as normal.
- People with **diabetes** should have additional instructions (contact the department on 01223 216546 immediately if missing).
- If you have any queries about the procedure or find that you cannot keep this appointment, please contact the Endoscopy Office between 0900 and 1700 on Tel: 01223 216546.

On the day

- Have **nothing to eat or drink for six hours** before your appointment
- If you have heart valve disease or require antibiotics when you visit the dentist please tell us when you come for the procedure.

At the hospital

- Please come to the Endoscopy Department, which is on Level 3 of the Addenbrooke's Treatment Centre (ATC).
- Use **Car park F**, situated opposite the main entrance to the centre, take the 'visitor' parking ticket issued at the car park entrance to the reception desk in the Treatment Centre to have it changed to the concession for outpatients' version. Bring your appointment letter with you.
- Please note that the appointment time is for your pre procedure check, not the time of your examination. You may be here for up to 2 hours.

Your doctor has requested this procedure to help investigate your medical condition to aid your diagnosis and management.

What is an ERCP?

ERCP stands for 'endoscopic retrograde cholangio-pancreatography'. It is a procedure that allows the endoscopist to examine the tubes that drain bile from your liver and gall bladder and digestive juices from the pancreas.

Bile is made in the liver, which is in the upper right part of the abdomen. Bile passes from liver cells into tiny tubes called bile ducts, these join together like the branches of a tree. Bile constantly drips down the bile duct into the duodenum (the first part of the gut after the stomach). Bile helps to digest food, particularly fatty foods.

The gallbladder lies under the liver on the right side of the upper abdomen. It is like a pouch which comes off the bile duct. It is a 'reservoir' which stores bile between meals. It contracts (squeezes) when you eat, emptying stored bile back into the bile duct.

The pancreas is a large gland that makes enzymes (chemicals), these enzymes flow into the duodenum. The pancreatic enzymes are vital to digest food.

Jaundice, which is yellowing of the skin and urine, occurs when the tubes draining the bile become blocked. ERCP procedures are undertaken to relieve this condition.

A duodenoscope is used, this is a flexible tube thinner than your index finger, with a light at the end. It is passed into the mouth, through the stomach to the duodenum to find the small opening (called Ampulla of Vater) where the bile and digestive juices drain into the intestine. A tube is passed through the duodenoscope and up into the Ampulla so that contrast (dye that can be seen on an x-ray) can be injected. X-rays are then taken.

Sometimes we need to make a small cut in the Ampulla so that gallstones which are stuck in the bile duct can be removed, this is called a sphincterotomy and is painless.

In other cases we need to put a tube called a stent into an area where the bile duct is blocked to allow the bile to drain.

Sometimes it is helpful to take a biopsy - sample of the lining of the ducts. This is done by passing a small instrument called forceps through the duodenoscope to 'pinch' out a tiny bit of the lining (about the size of a pinhead) which is sent to the laboratory for analysis.

These procedures usually take about 15 minutes but times vary considerably. If it takes longer, you should not worry.

Getting ready for the procedure

On arrival, the procedure will be explained and you will be asked to sign a consent form. We want to make sure that you understand the procedures and their implications. Remember, you can change your mind about having the procedure at any time.

You will be given an intravenous sedative (this is **not** a general anaesthetic). It is an injection into a vein, which will make you feel relaxed and sleepy but not unconscious. Some people will also need to have antibiotics.

The sedative will continue to have a mild effect for up to 24 hours and may leave you unsteady on your feet for a while.

You **must** arrange for a responsible adult to come with you, wait with you and then take you home. Altogether, you and your escort may be in the department for up to two hours.

If you come without someone, the procedure will be cancelled. If you are using hospital transport, an escort is not required.

Please leave any valuables at home; you will need to undress and put on a gown.

During the procedure

For your comfort and reassurance, a trained nurse will stay with you throughout. In the procedure room, you will be asked to remove false teeth and glass, and made comfortable on the x-ray table lying on your left side with your left arm behind your back.

The endoscopist will give you the sedative injection. So that you do not bite the duodenoscope during the procedure, a plastic mouth guard will be put gently between your teeth. You will be given oxygen through a facemask, and a plastic 'peg' will be placed on your finger to monitor your pulse and oxygen levels.

As the tube is gently passed through your mouth you may gag slightly, this is quite normal and will not interfere with your breathing. During the procedure some air will be put in to your stomach so that the endoscopist will have a clear view. This may make you burp and belch a little, some people find this uncomfortable. The air is removed at the end and when the procedure is finished, the duodenoscope is removed quickly and easily.

Minimal restraint may be appropriate during the procedure. However if you make it clear that you are too uncomfortable the procedure will be stopped.

Potential problems

This procedure involves x-rays; **if you are pregnant, you should not have this procedure.**

ERCP procedures carry a very small risk (5 in 1000 cases) of haemorrhage (bleeding) or perforation (tear).

If a cut is made into the bile duct there is a risk of 1 in 50 of significant bleeding. This can be treated straight away through the duodenoscope and rarely is a major complication, however if it is severe sometimes blood transfusion or surgery is needed.

Occasionally inflammation of the pancreas (pancreatitis) may develop (1 in 50 – 100), it can be painful and usually requires you to stay in hospital for a few more days for intravenous fluids and painkillers. On very rare occasions, it may be more severe than this.

There may be a slight risk to crowned teeth or dental bridgework and you should tell the endoscopist if you have any of these.

Another rare complication is an adverse reaction to the intravenous sedative and antibiotic drugs.

You can be reassured that your doctors will only have recommended ERCP if the benefit to you from the procedure clearly outweighs these small risks.

After the procedure

Following the procedure, you will be taken to a recovery area while the sedation wears off. When you are sufficiently awake, you will be given a drink before getting dressed. You can then go home; this may be up to an hour following the procedure.

The back of your throat may feel sore for the rest of the day. You may also feel bloated if some air remains in your stomach. Both these discomforts will pass and need no medication.

You are advised not to drive, operate machinery, return to work, drink alcohol or sign any legally binding documents for the next 24 hours. You are also advised to have a responsible adult to stay with you for the next 12 hours.

If you have any of the following you should contact your GP, the Endoscopy Department or the Accident and Emergency Department for further advice.

- severe pain,
- black tarry stools
- persistent bleeding

When do I know the result?

The endoscopist will be able to tell you the results immediately after the procedure. It is a good idea to have someone with you when you talk to the endoscopist because the sedation can make you forget what has been discussed.

If biopsies were taken you will be told the final diagnosis by the team who requested the ERCP (in the clinic or by letter to you or your GP). These results may take several weeks to come through.

Details of the results and any further treatment should be discussed with the doctor who recommended you have this procedure.

Training doctors and other health professionals is essential to the continuation of the National Health Service, and improving the quality of care. Your treatment may provide an important opportunity for such training under the careful supervision of a senior doctor.

You can, however, decline to be involved in the formal training of medical and other students: this won't affect your care and treatment.

Alternatives:

There are no real alternatives to therapeutic ERCP. In some cases, depending on individual factors such as the symptoms present and the condition being investigated, alternatives may be:

- MRCP or PTC.

For more information:

- Contact the Endoscopy Office between 0900 and 1700 on Tel: 01223 216546.
- See www.addenbrookes.org.uk/consent

Please ask if you require this information in other languages, large print or audio format: 01223 216032 or patient.information@addenbrookes.nhs.uk

Potete chiedere di ottenere queste informazioni in altre lingue, in stampato grande o in audiocassetta.

Italian

若你需要本信息的繁體中文、大字體或音訊格式的版本，請要求索取。

Cantonese

તમને આ માહિતી બીજી ભાષાઓમાં, મોટા અક્ષરોમાં અથવા સાંભળી શકાય એવા માધ્યમ (ઓડીઓ ફોર્મેટ)માં જોઈતી હોય તો કૃપા કરીને પૂછો.

Gujarati

تکایہ پرسیار بکہ نہ گہر نہ وزانیاریہت دہوی بہ زمانیکی تر , بہ پیتی گہورہ یانیش بہ شیوہی دہنگ

Kurdish

اگر آپ کو یہ معلومات دوسری زبانوں میں، بڑے الفاظ کی اشاعت میں یا آڈیو ٹیپ پر درکار ہوں تو براہ مہربانی اس کیلئے درخواست کریں۔

Urdu



Addenbrooke's is smoke-free. Please do not smoke anywhere on the site.

For advice on quitting, contact your GP or the NHS smoking helpline free, 0800 169 0 169

Document history

Authors	Q Tuffnell, A Wright
Department	Endoscopy Department, Box 293, Addenbrooke's Hospital, Cambridge University Hospitals NHS Foundation Trust, Hills Road, Cambridge, CB2 0QQ www.addenbrookes.org.uk
Contact number	01223 216515
Published	June 2007
Review date	June 2009
File name	Endo_ercp_jo.doc (word)
Version number	1
Ref	655