
Ophthalmology

Fluorescein Angiography

Fluorescein Angiography is a routine retinal diagnostic procedure, which gives information on the retinal and optic nerve blood vessels at the back of the eye. It is particularly useful in diabetes but it is also helpful in many other eye disorders.

Before the procedure

Before the procedure, you will have drops put in your eyes to enlarge the pupils. The drops take from 20 to 30 minutes to fully dilate your pupils allowing the photographer to view the back of your eye.

These drops will cause temporary blurring of your near vision, possibly your distance vision and sensitivity to bright lights. This usually lasts about four hours but can last up to 24 hours. It is therefore advisable to make appropriate transport arrangements. Bring sunglasses with you if you have them.

The procedure

The procedure is performed while you are seated with your chin on a rest and your forehead against a headband. It is important that you are comfortable for the procedure.

A doctor or specially trained nurse injects a small quantity of fluorescein dye into a vein in your arm or hand. The camera flashes repeatedly to take the photographs. Please follow the photographer's instructions as closely as you can. It is important not to move your head or eyes unless you are told to do so. The procedure lasts about ten minutes.

Side effects of the treatment

- Fluorescein angiography involves injecting an intravenous dye. There is a remote risk of life threatening, adverse reaction requiring immediate treatment. The procedure is, therefore, always performed with emergency resuscitation facilities and trained medical staff immediately available.

- The fluorescein dye will give your skin a yellowish tinge and will also turn your urine bright yellow for a few hours following the procedure.
- You may develop a transient rash or itch.
- Some patients feel nauseated during the test. If you take a few deep breaths, the nausea usually passes rapidly.
- There should be no discomfort from the procedure and care is taken to ensure that the dye flows freely. Very occasionally the dye can leak around the injection site causing pain for a few hours.
- The risk of death from fluorescein angiographic injections is less than 1 in 200,000.
- Serious reactions from fluorescein angiography are extremely rare, except in patients with severe asthma or allergic diseases. Please tell the doctor if any of the following apply to you:
 - Asthma
 - Allergy to any medicines, food or animals
 - Taking any medication
 - Previous reaction to fluorescein
 - Pregnancy

Please ask if you require this information in other languages, large print or audio format: 01223 216032 or patient.information@addenbrookes.nhs.uk

Potete chiedere di ottenere queste informazioni in altre lingue, in stampato grande o in audiocassetta.

Italian

若你需要本信息的繁體中文、大字體或音訊格式的版本，請要求索取。

Cantonese

तमने आ माहिती वीछ भाषाओमां, मोटा अक्षरोमां अथवा सांभली शकाय जेवा माध्यम (ओडीओ इमेज)मां जेठती छीय तो कृपा करीने पूछो.

Gujarati

تکایہ پرسیار بکے نہ گھر نہ وزانیاریہت دہوی بہ زمانیکی تر , بہ بیٹی گہورہ یانیش بہ شیوہی دہنگ

Kurdish

آگر آپ کو یہ معلومات دوسری زبانوں میں، بڑے الفاظ کی اشاعت میں یا آڈیو ٹیپ پر درکار ہوں تو براۓ مہربانی اس کیلئے درخواست کریں۔

Urdu



Addenbrooke's is smoke-free. Please do not smoke anywhere on the site.
For advice on quitting, contact your GP or the NHS smoking helpline free, 0800 169 0 169

Document history

Authors	Mr Flanagan
Department	Department of Ophthalmology, Addenbrooke's Hospital, Cambridge University Hospitals NHS Foundation Trust, Hills Road, Cambridge, CB2 2QQ www.addenbrookes.org.uk
Contact number	01223 216577
Published	January 2007
Review date	January 2009
File name	Fluorescein_angiography.doc
Version number	1
Ref	PIN 615