
Department of Oral and Maxillofacial Surgery

Frequently Asked Questions: Removal of Impacted Wisdom Teeth

This leaflet has been written to help your understanding of the removal of wisdom teeth and answers many commonly asked questions. If you have any other questions that the leaflet does not answer or would like further explanation please ask your surgeon. This leaflet has been taken from the British Association of Oral and Maxillofacial Surgeons.

The problem

The wisdom teeth (third molars) are the last teeth to grow. They grow into the mouth, through the gums, anytime after about 16 years of age. Frequently there is not enough room in the mouth for wisdom teeth and so they do not grow into the mouth normally. When this happens, the wisdom teeth are said to be “impacted”. Wisdom teeth are usually either impacted forwards into the tooth in front or backwards into the jaw bone.

Why do I need treatment?

An impacted wisdom tooth can cause a number of problems that mean the tooth is best removed. Most commonly these are:

- Repeated attacks of infection in the gum surrounding the tooth. This leads to pain and swelling.
- Food packing, which causes decay in either the wisdom tooth or the tooth in front.
- Cysts can form around the wisdom tooth if it does not come into the mouth properly. A cyst occurs when fluid fills the sack that normally surrounds a developing wisdom tooth.

What does the treatment involve?

- Because the wisdom tooth has not fully grown into the mouth it is often necessary to make a cut in the gum over the tooth.
- Sometimes it is also necessary to remove some bone surrounding the wisdom tooth to allow the tooth to be removed.
- Sometimes the tooth will need to be cut into two or three pieces to make it easier to remove.
- Once the wisdom tooth has been removed, the gum is put back into place with stitches.
- In the majority of cases these stitches are dissolvable and take around two weeks to disappear.

What type of anaesthetic is used?

A number of options are available and depend on how difficult the wisdom tooth is to remove:

- **Local anaesthetic:** this is an injection into the gum surrounding the wisdom tooth, rather similar to what you may have had at your dentist for a filling. The injection takes a couple of minutes to numb the area and means that you will feel no pain while the wisdom tooth is removed. This is the best option for wisdom teeth that are simple to remove.
- **Local anaesthetic and intravenous sedation:** in addition to a local anaesthetic injection you can be given an injection into your arm. This makes you feel relaxed and less aware of the procedure.
- **General anaesthetic:** it is usually possible to remove wisdom teeth under a “day case” general anaesthetic. This means that although you are put to sleep completely, you will be able to go home on the same day as surgery.

How long does it take to remove a wisdom tooth?

This varies; some wisdom teeth may take only a few minutes to remove. More difficult wisdom teeth that need to be cut into pieces to remove can take around 20 minutes to remove.

Is there much pain or swelling after the removal of wisdom teeth?

It is likely that there will be some discomfort and swelling both on the inside and outside of your mouth after surgery. This is usually worse for the first three days, but it may take up to two weeks before all the soreness goes. You may also find that your jaw is stiff and you may need to eat a soft diet for a week or so. If it is likely to be sore your surgeon will arrange painkillers for you. It may also be necessary for you to have a course of antibiotics after the extraction. There may be some bruising of the skin of your face that can take up to a fortnight to fade away.

Is there anything else I need to do after the extractions?

It is important to keep the extraction sites as clean as possible for the first few weeks after surgery. It may be difficult to clean your teeth around the sites of the extraction because it is sore and if this is the case it is best to keep the area free from food debris by gently rinsing with a mouthwash or warm salt water (dissolve a flat teaspoon of kitchen salt in a cup of warm water) commencing on the day after surgery.

Do I need to take any time off work?

Usually it will be necessary to take the next day off work to recover from the surgery, and avoid strenuous exercise for 24 hours as this may cause bleeding. Depending on the type of anaesthetic used you may well not be able to drive (12 hours after intravenous sedation and for 24 hours after a general anaesthetic).

What are the possible problems?

- Although there may be a little bleeding at the time of the removal, this usually stops very quickly and is unlikely to be a problem if the wound is stitched. Should the area bleed again when you get home this can usually be stopped by applying pressure over the area for at least 10 minutes with a rolled up handkerchief or swab. If the bleeding does not stop, please contact the department.
- Infection is uncommon particularly if the mouth is kept clean.
- There are two nerves that lie very close to the roots of the lower wisdom teeth.
 - One of these nerves supplies feeling to your lower lip, chin and lower teeth. The other supplies feeling to your tongue and helps with taste.
 - Sometimes these nerves may be bruised when a wisdom tooth is taken out.
 - This can cause tingling or numbness in your lip, chin or tongue, and more rarely altered taste.
 - About one in 10 people will have some tingling or numbness that can last several weeks.
 - Less than one in 100 people will have problems that last more than a year.
 - These risks may be higher if your tooth is in a difficult position. Your surgeon will tell you if you are considered to be at an increased risk.

What are the alternatives to this treatment?

There are no surgical or non-surgical alternatives to this treatment.

After surgery

After the removal of wisdom teeth it is not usually necessary to return for a follow up appointment. The stitches dissolve. If there are problems, contact the department.

For more information:

Department of Oral and Maxillofacial Surgery

Addenbrooke's Hospital, Cambridge University Hospitals NHS Foundation Trust, Hills Road, Cambridge, CB2 0QQ

Tel: 01223 216635

Website: www.addenbrookes.org.uk/omfs

NICE Guidelines for wisdom teeth

www.nice.org.uk

Adapted from:

British Association of Oral and Maxillofacial Surgeons

<http://www.baoms.org.uk/>

Please ask if you require this information in other languages, large print or audio format:
01223 216032 or patient.information@addenbrookes.nhs.uk

Potete chiedere di ottenere queste informazioni in altre lingue, in stampato grande o in audiocassetta.

Italian

若你需要本信息的繁體中文、大字體或音訊格式的版本，請要求索取。

Cantonese

तमने आ माळिती बीछ भाषाओमां, मोटा अक्षरोमां अथवा सांभजी शकाय जेवा माध्यम
(ओडीओ डोमेट)मां जेठती छोय तो कृपा करीने पूछो.

Gujarati

تکایہ پرسیار بگہ نہ گہر نہ وزانیاریہت دہوی بہ زمانیکی تر . بہ پیتی گہورہ یانیش بہ شیوہی دہنگ

Kurdish

آگراپ کو یہ معلومات دوسری زبانوں میں، بڑے الفاظ کی اشاعت میں یا آڈیو ٹیپ پروکاروں تو ہر اے مہربانی اس کیلئے درخواست کریں۔

Urdu



Addenbrooke's is smoke-free. Please do not smoke anywhere on the site.
For advice on quitting, contact your GP or the NHS smoking helpline free, 0800
169 0 169

Document history

Authors	David Adlam
Department	Oral and Maxillofacial Surgery Addenbrooke's Hospital, Cambridge University Hospitals NHS Foundation Trust, Hills Road, Cambridge, CB2 0QQ www.addenbrookes.nhs.uk
Contact number	01223 216635
Published	October 2005
Reviewed and reissued	August 2007 (no changes made)
Review date	August 2009
File name	FAQS_Wisdom_Teeth.doc
Version number	2
Ref	PIN482